

Selection of compost-derived actinomycetes with plant-growth promoting and tomato stem rot biocontrol potentialities

Fakher Ayed^{1,2,3*}, Rania Aydi Ben Abdallah³, Hayfa Jabnoun-Khiareddine³, Mejda Daami-Remadi³

¹National Agronomic Institute of Tunisia, 1082 Tunis, University of Carthage, Tunisia

²Technical Centre of Organic Agriculture, 4042 Chott-Meriam, Sousse, Tunisia

³UR13AGR09-Integrated Horticultural Production in the Tunisian Centre-East, Regional Research Centre on Horticulture and Organic Agriculture, University of Sousse, 4042, Chott-Meriem, Tunisia

Abstract

Seventeen actinomycetes isolates, recovered from 2 composts, were screened for their ability to promote the growth of tomato seedlings and to suppress stem rot disease caused by *Sclerotium rolfii*. Tomato cv. Rio Grande seedlings inoculated with *S. rolfii* and treated with A2-3, A3-3, A4-3, A5-3, A8-3, A9-3, A1-4, A2-4, A3-4, A4-4, A6-4, and A10-4 actinobacterial isolates showed 23.3-70% less disease severity than the inoculated and untreated controls. A3-3, A2-4, and A4-4 based treatments applied to *S. rolfii*-infected tomato seedlings had significantly enhanced all growth parameters as compared to control. The recorded increments were estimated at 35.52-66.6% for height, 37.4-53.4% for the stem diameter, 38.5-95.6% for the aerial part dry weight, and 81.8-151% for the root dry weight. Treatments with A3-3 and A4-4 isolates had increased the majority of tomato growth parameters by 15.8-56.5% over the pathogen-free control. Tomato seedlings treated with A4-3 and A1-4 isolates showed between 35.2-22.8% and 42.3-43.3% higher aerial part dry weight and root dry weight, respectively, as compared to pathogen-free and untreated control. This investigation demonstrated that the tested composts can be explored as potential sources for the isolation of actinomycetes acting as biocontrol and bio-fertilizing agents.

Keywords: actinobacteria, antifungal activity, growth enhancement, *Sclerotium rolfii*, *Solanum lycopersicum*.

*Corresponding author: Fakher Ayed,
E-mail: ayedfakher@yahoo.fr

1. Introduction

Composting is defined as a biochemical process managed under principally thermophilic and aerobic conditions by active microorganisms to produce a renewable organic resource (Bohacz, 2018). This process is a sustainable option for recycling various agro-industrial wastes, on small or large scale, to obtain a mature and stable organic matter (Scotti et al., 2016). The final product, *i.e.*, compost, can be used to improve soil physicochemical and microbiological properties and suppress various soilborne diseases (Pane et al., 2019; Stavi et al., 2016). These effects are partly attributed to their associated beneficial microorganisms and bioactive metabolites (Hadar & Papadopoulou, 2012). Thus, composts are becoming an alternative tool for the extensive use of synthetic inputs in cropping systems and are deemed safer for the environment and human health (Coventry et al., 2006). In the past decades, the added value of composts attributed to their disease-suppressive effects have been extensively demonstrated against various soilborne pathogens such as *Rhizoctonia solani*, *Verticillium dahliae*, *Fusarium* spp., *Sclerotinia* spp., *Phytophthora* spp., *Pythium* spp., and *Thielaviopsis* sp. causing plant wilting, damping-off and decaying in many important crops (Tubeileh & Stephenson, 2020; Pane et al., 2013; Alfano et al., 2011; Bonanomi et al., 2007). The compost disease-suppressive potential is mainly related to the biological activity of its associated microbiota, which interacts with the soil organic matter and the host plant by regulating the microbial communities in the rhizosphere (Hadar & Papadopoulou, 2012; Manici et al., 2004), and to its capacity to improve plant nutrition and growth (Martin, 2015). Among the widely documented compost-associated microbial agents, bacteria, actinomycetes, and fungi were the focal driving forces for plant diseases suppression (Coelho et al., 2020; Joshi et al., 2009). These microbial agents acted against target pathogens through antibiosis,

hyperparasitism, and competition for space and nutrients (Larkin & Tavantzis, 2013). Actinomycetes, gram-positive bacteria, and members of the *Actinobacteria* group are well known as secondary metabolites producers and are widely explored for various agricultural features (Qin et al., 2011). *Actinomycetes* have been recovered from diverse natural environments such as rhizospheric soil, composts, and healthy plant tissues. Commercial bio-molecules are mostly produced by *Streptomyces*, *Saccharopolyspora*, *Micromonospora*, *Amycolatopsis*, and *Actinoplanes* (Palla et al., 2018). These microbial agents can improve plant growth and support its establishment even under stress conditions (Srivastava et al., 2015; Hamdali et al., 2008). Moreover, their antagonistic potential against phyto-pathogenic organisms was demonstrated (Nurkanto & Julistiono, 2014; Anouar et al., 2012). They can protect roots by inhibiting the fungal pathogen development mostly through the production of antifungal compounds or cell wall degrading enzymes (Bhatti et al., 2017). *Sclerotium rolfsii* Sacc. is a soilborne pathogen that affects a wide host range of over 500 dicotyledonous and monocotyledonous plant species (Sun et al., 2020; Punja, 1985; Aycok, 1966). Infection by this pathogen may occur at all growing stages and lower stems at or near the soil surface by forming water-soaked lesions. These lesions spread quickly to girdle stems where white mycelial mats may be observed on the infected plant tissues. Severely infected plants may wilt thus leading to partial or total yield loss (Sun et al., 2020; Kator et al., 2015; Fery & Dukes, 2002). Stem rot diseases are usually managed using chemical fungicides such as carboxin, carbendazim, benomyl, propiconazole, methyl thiophanate, and oxycarboxin (Sridharan et al., 2020). Nevertheless, the hazardous use of these chemicals represents a severe threat to the environment, food safety, and human health (Sridharan et al., 2020). Biological control of soilborne phytopathogens is of increased interest where various microorganisms

recovered from diverse ecological niches are largely explored for sustainable management of stem rot disease (Singh et al., 2013). Among the explored microorganisms, actinomycetes are considered as promising potential candidates for the suppression of *S. rolfsii* (Anusree & Bhai, 2017). Therefore, the main objectives of this investigation are: (1) to isolate actinomycetes associated with two selected composts; (2) to evaluate their ability to promote the growth of tomato seedlings and to control stem rot disease.

2. Materials and methods

2.1 Pathogen growth conditions

S. rolfsii isolate Sr₁, used in the current study, was originally recovered from potato plants showing typical stem rot symptoms. Identification and pathogenicity of this isolate were previously investigated (Ayed et al., 2018a; Daami-Remadi et al., 2010). The pathogen was grown on Potato Dextrose Agar (PDA) medium for 15 days at 30 °C, in the dark, before used.

2.2 Composts preparation

Bioassays were carried out using 2 mature composts. The first one (C₃) are composed by a 70% of cattle and 25% of chicken manures mixture associated with 5% green waste, the second one (C₄) are issued from 70% of cattle and 25% of sheep manures mixed with 5% of olive-mill solid waste. The composting system was carried out in parallel open-windrows in the following dimensions: height 1.5 m × base width 2.0 m × length 10 m. The compost mass was mechanically homogenized and the humidity of the mass during fermentation was maintained using water at an optimal level (60-70%) for the composting process for 8 months. These two locally produced composts and their teas were screened for their plant growth-

promoting potential and for their ability to control stem rot disease on tomato seedlings (Ayed et al., 2018b; 2018c).

2.3 Isolation of compost-associated actinomycetes

A sample of 10 g of air-dried compost was suspended into a 90 ml volume of sterile distilled water (SDW) in a 250 ml flask, and shaken for 30 min at 200 rpm. Serial dilution of compost was carried out, and then a sample of 100 µl was spread onto Petri-dish containing Actinomycetes Isolation Agar medium amended with 500 µg/ml of streptomycin sulfate, 100 µg/ml of chloramphenicol, 50 µg/ml of ampicillin, and 100 µg/ml of cycloheximide (Joshi et al., 2009). Plates were maintained at 28 ± 2 °C for 14-21 days, in darkness. Developed colonies showing distinct macro-morphological traits were isolated separately on Yeast Malt Agar (ISP-2), and stored in glycerol (20%) at -20 °C for further use (Himaman et al., 2016).

2.4 Morphological characterization of the actinomycete isolates

Obtained isolates were grown in ISP-2 medium at 28 ± 2° C for 10 days. Actinobacterial colonies were characterized based on the colony appearance, the type of areal hyphae, and the growth of vegetative hyphae. The color of diffusible pigment production was visually estimated with the help of RHS-color code (RHS color chart, Fifth Edition-Royal Horticultural Society) (Anusree & Bhai, 2017).

2.5 Screening for actinomycetes growth-promoting ability

Actinomycete isolates were assessed for their ability to promote the tomato seedlings growth. Tomato cv. Rio Grande seeds were

surface-sterilized by immersing in 5% of sodium hypochlorite (NaOCl) for 3 min, then rinsed 6 times with SDW and air-dried (Aydi Ben Abdallah et al., 2018). Disinfected seeds were soaked for 30 min into actinobacterial cell suspensions ($\sim 10^8$ CFU/ml), prepared in ISP-2 liquid cultures incubated at 28 ± 2 °C for 48 h. They were sown in alveolated trays (4 cm \times 4 cm \times 4 cm) filled with commercialized peat (Klasmann-Deilmann, Bio-Substrate, Germany), and watered regularly with tap water. A volume of 10 ml of each actinobacterial suspension was applied per alveolus by substrate drench immediately, 15 days, and 30 days post-sowing. Controls were soaked in SDW and watered later regularly with tap water. Ten seedlings were used for each treatment. The bioassay was carried out under plastic tunnels at 26-32 °C with 12 h (light) / 12 h (dark) photoperiod and 60% air relative humidity. At 35 days post-sowing, tomato seedlings were uprooted and washed for removing adhering peat. Different growth parameters were recorded (plant height, stem diameter, aerial part, and root dry weights). Data were analyzed according to a completely randomized design. The whole experiment was repeated twice.

2.6 Screening for actinomycetes stem rot suppression ability

Tomato cv. Rio Grande seeds, disinfected as detailed above, were soaked for 30 min into a water cell suspension (10^8 CFU/ml) of collected actinomycetes isolates and sown in alveolated trays filled with commercialized peat (Aydi Ben Abdallah et al., 2018). All tested treatments were applied directly, 15 days and 30 days post-sowing (10 ml per alveolus per individual treatment). Challenge inoculation with *S. rolf sii* was performed as substrate drench with 20 ml of a mixture of mycelium and sclerotia (≈ 25 -30 sclerotia per

100 ml) 21 days post-sowing (Daami-Remadi et al., 2012). Uninoculated and inoculated controls were watered with the same volume of SDW and watered later regularly with tap water. The bioassay was carried out using ten seedlings per individual treatment and maintained under the same greenhouse conditions as indicated above. At 35 days post-sowing, disease severity was recorded based on a 1-5 scale (De Curtis et al., 2010), where 1 = no stem lesion, 2 = lesions girdled $\leq 25\%$ of the stem circumference, 3 = lesions girdled 26-50% of the stem circumference, 4 = lesions girdled $> 51\%$ of the stem circumference and 5 = stem completely girdled. Plant height, stem diameter, root, and aerial part dry weights were also noted. Data were analyzed according to a completely randomized design. The whole experiment was repeated twice.

2.7 Statistical analysis

The data were statistically analyzed by analysis of variance (ANOVA) with the statistical package SPSS software (Version 20) and subjected to mean separation by Duncan Multiple Range test (at $P \leq 0.05$). Correlations between disease severity and seedling growth parameters were performed using bivariate Pearson's test at $P \leq 0.05$.

3. Results

3.1 Isolation of compost-associated actinomycetes and their morphological characterization

A collection of 17 actinomycetes isolates, exhibiting diversity in their macro-morphological traits, was recovered from the 2 composts C₃ and C₄. The color of aerial mycelium of actinomycetes isolates varied from white to brownish-black, whereas the

color of submerged mycelium varied from white to black. Colonies were circular or irregular, raised or flat or punctiform, and entire or undulate (Table 1).

Table 1: Cultural characteristics of actinomycetes isolates on ISP-2 medium after incubation at $28 \pm 2^\circ\text{C}$ for 10 days.

Isolates	Composts	Color of aerial mycelium	Color of submerged mycelium	Colonies characteristics
A1-3	C ₃	White	White	Circular, raised, entire
A2-3	C ₃	Brownish-black	Brown	Circular, flat, entire
A3-3	C ₃	Brownish-black	Brown	Irregular, raised, undulate
A4-3	C ₃	Brownish-black	Brown	Circular, flat, punctiform
A5-3	C ₃	Greyed-brown	Brown	Irregular, raised, undulate
A 8-3	C ₃	Greyed-white	White	Circular, raised, entire
A9-3	C ₃	White	White	Circular, flat, punctiform
A10-3	C ₃	Greyed-black	Black	Circular, punctiform, raised
A1-4	C ₄	White	White	Circular, punctiform, entire
A2-4	C ₄	White	Greyed-yellow	Circular, flat, entire
A3-4	C ₄	Greyed-brown	Greyed-brown	Irregular, raised, undulate
A4-4	C ₄	Greyed-brown	Greyed-brown	Irregular, raised, undulate
A5-4	C ₄	White	Greyed -white	Circular, raised, entire
A6-4	C ₄	Greyed-green	Green	Circular, flat, entire
A7-4	C ₄	White	White	Circular, raised, entire
A8-4	C ₄	Greyed-white	White	Circular, raised, entire
A10-4	C ₄	Grey	Greyed-white	Circular, raised, entire

3.2 Disease suppression ability of compost-associated actinomycetes

ANOVA analysis revealed that stem rot severity, noted after 35 days post-sowing and 14 days post-inoculation, varied significantly depending on the tested actinobacterial treatments. As shown in Figure (1), a significant decrease in disease severity by 23.33 to 70% over pathogen-inoculated and untreated control was noted on tomato seedlings infected with *S. rolf sii* and treated with A2-3, A3-3, A4-3, A5-3, A8-3, A9-3, A1-

4, A2-4, A3-4, A4-4, A6-4, and A10-4 isolates. Tomato seedlings inoculated by the pathogen and treated with these isolates showed a significantly disease severity similar to the pathogen-free and untreated ones. The highest decreases in stem rot severity, of about 56.67 and 70%, were noted on tomato seedlings treated with A9-3 and A4-4 isolates, respectively. However, no disease-suppressive effect, as compared to the inoculated and untreated control, was noted on those treated with A1-3, A10-3, A5-4, A7-4, and A8-4 isolates.




Figure 1: Effect of compost-associated actinomycetes isolates on stem rot severity noted after 35 days post-sowing on *Sclerotium rolf sii*-inoculated tomato cv. Rio Grande seedlings as compared to the untreated controls. UC: Uninoculated and untreated control; IC: *S. rolf sii*-inoculated and untreated control. Results are presented as means \pm SE (n = 10, $P \leq 0.05$). Bars sharing the same letter are not significantly different according to Duncan's Multiple Range test (at $P \leq 0.05$).

3.3 Growth-promoting ability of compost-associated actinomycetes on *Sclerotium rolfii*-inoculated tomato seedlings

Growth parameters of tomato seedlings, noted after 35 days post-sowing and 14 days post-inoculation with *S. rolfii*, varied significantly (at $P \leq 0.05$) depending on tested biological treatments. As illustrated in Table (2), treatments with A2-3, A3-3, A4-3, A5-3, A8-3, A9-3, A10-3, A1-4, A2-4, A3-4, A4-4, A5-4-, and A6-4 isolates significantly improved the seedling height by 21.1 to 66.6% over

pathogen-inoculated and untreated control, where the highest increment was induced by A4-4 treatment. Furthermore, a significant enhancement in the stem diameter, by 37.4 to 62.2% over pathogen-inoculated control, was recorded on tomato seedlings treated with A2-3, A3-3, A2-4, A3-4, and A4-4 isolates. A3-3 and A4-4 based treatments exhibited the highest growth-promoting effect based on the aerial part dry weight which was significantly improved by 95.6% and 66.2%, respectively, over the pathogen-inoculated and untreated control.

Table 2: Growth-promoting potential of compost-associated actinomycete isolates noted 35 days post-sowing on tomato cv. Rio Grande seedlings inoculated with *Sclerotium rolfii* as compared to the untreated controls.

Biological treatments	Plant height (mm)	Stem diameter (mm)	Aerial part dry weight (mg)	Root dry weight (mg)
UC	81.5 ± 1.35 b	2.28 ± 0.1 a	189.87 ± 7.10 bcdef	26 ± 1.13 cdefg
IC	59.12 ± 1.17 de	1.38 ± 0.11 ef	142.25 ± 13.91 fg	17.87 ± 1.25 fgh
A1-3	39.75 ± 5.28 f	0.44 ± 0.24 g	58.37 ± 23.03 h	11 ± 1.88 h
A2-3	73 ± 6.16 bc	2.24 ± 0.23 a	145.75 ± 8.81 efg	18.37 ± 0.71 fgh
A3-3	80.12 ± 6.96 b	2.03 ± 0.21 abc	278.25 ± 17.78 a	39.37 ± 5.08 ab
A4-3	82.25 ± 5.43 b	1.77 ± 0.16 bcde	214.62 ± 17.86 bc	36.37 ± 4.97 abc
A5-3	81.75 ± 2.39 b	1.57 ± 0.11 cde	176.75 ± 10.71 cdef	24.37 ± 0.98 defg
A8-3	73.12 ± 2.53 bc	1.78 ± 0.13 bcde	184.75 ± 17.60 cdef	30.62 ± 6.61 bcde
A9-3	72.37 ± 2.8 bc	1.65 ± 0.66 cde	146.5 ± 11.25 efg	19.75 ± 0.90 efgh
A10-3	73 ± 2.1 bc	1.76 ± 0.14 bcde	169.37 ± 15.82 cdefg	25 ± 1.80 defg
A1-4	76.37 ± 5.53 bc	1.65 ± 0.20 cde	202.5 ± 16.86 bcd	27.87 ± 3.60 cdef
A2-4	80.5 ± 3.9 b	2.12 ± 0.08 ab	197 ± 11.27 bcd	32.5 ± 3.02 bcd
A3-4	80.37 ± 4.23 b	2.02 ± 0.09 abc	208.12 ± 10.34 bc	24 ± 2.15 defg
A4-4	98.5 ± 3.57 a	1.9 ± 0.07 abcd	236.37 ± 18.32 ab	44.87 ± 6.16 a
A5-4	73.62 ± 3.8 bc	1.63 ± 0.08 cde	195 ± 20.54 bcde	36.25 ± 6.39 abc
A6-4	71.62 ± 1.77 bc	1.69 ± 0.09 bcde	184.37 ± 3.73 cdef	27.37 ± 2.07 cdef
A7-4	49.12 ± 0.55 ef	1.14 ± 0.08 f	140.37 ± 4.66 fg	22 ± 2.36 defgh
A8-4	64.62 ± 4.5 cd	1.53 ± 0.19 def	127.75 ± 25.73 g	16.37 ± 3.13 fgh
A10-4	50.87 ± 2.71 ef	0.34 ± 0.03 g	153.37 ± 5.55 defg	14.62 ± 1.89 gh

UC: Uninoculated and untreated control; IC: *S. rolfii*-inoculated and untreated control. Results are presented as means ± SE (n = 10, $P \leq 0.05$). For each column, values followed by the same letter are not significantly different according to Duncan's Multiple Range test (at $P \leq 0.05$).

Treatments with A4-3, A1-4, A2-4, A3-4, and A5-4 isolates led to a significant increment in this parameter by 37.1-50.9%. As for their effects on the root dry weight, A3-3, A4-3, A4-4, and A5-4 isolates had significantly improved this parameter by 102.8-151.1% over pathogen-inoculated and untreated control. It should be also highlighted that A8-3 and A2-4 based treatments led to a 71.3% and 81.8% increase in the root dry weight over control, respectively. However, A1-3, A7-4, A8-4, and A10-4 isolates had no positive

effects based on all studied growth parameters.

3.4 Correlation between stem rot severity and seedling growth parameters

Pearson's correlation analysis revealed that the lowest stem rot severity, estimated based on a necrosis index, led to significant increases in all seedling growth parameters. Indeed, the seedling height was negatively correlated to necrosis index ($r = -0.471$; $n=190$; $P =$

0.000). Moreover, the stem diameter was negatively linked to stem rot severity ($r = -0.431$; $n=190$; $P = 0.000$). Furthermore, the aerial part dry weight ($r = -0.392$; $n=190$; $P = 0.000$) and the root dry weight ($r = -0.268$; $n=190$; $P = 0.001$) were negatively correlated to the necrosis index.

3.5 Growth-promoting ability of compost-associated actinomycetes on pathogen-free tomato seedlings

ANOVA analysis revealed that all growth parameters (height, stem diameter, aerial part dry weight, and root dry weight), varied significantly (at $P \leq 0.05$) depending on tested actinobacterial treatments. As presented in Table 3, the

A4-4 isolate exhibited the highest growth-promoting ability on tomato pathogen-free seedlings where the height was enhanced by 27% over the untreated control. This parameter was also significantly increased by 15.8 and 12.1% over control following seedling treatment with A3-3- and A3-4 isolates, respectively. However, no positive effect was noted based on the stem diameter. Treatment of tomato seedlings with A3-3, A4-3, A1-4, and A4-4 isolates resulted in a significant increment of the aerial part and root dry weights by 20.1-56.5% and 42.3-51%, respectively, over control. A3-3 and A5-4 isolates exhibited the highest growth-promoting potential based on the aerial part dry and the root dry weight, respectively (Table 3).

Table 3: Growth-promoting potential of compost-associated actinomycete isolates noted 35 days post-sowing on pathogen-free tomato cv. Rio Grande seedlings as compared to the untreated control.

Biological treatments	Plant height (mm)	Stem diameter (mm)	Aerial part dry weight (mg)	Root dry weight (mg)
UC	81.5 ± 1.35 def	2.28 ± 0.10 ab	189.87 ± 7.10 de	26 ± 1.13 bcdef
A1-3	69.37 ± 2.54 g	1.86 ± 0.06 bc	143.5 ± 11.10 f	19.37 ± 0.82 def
A2-3	81.37 ± 3.74 def	1.81 ± 0.01 bc	142.75 ± 1.38 f	18 ± 0.27 ef
A3-3	94.37 ± 2.56 b	2.43 ± 0.16 a	297.12 ± 23.67 a	39 ± 5.63 a
A4-3	90.25 ± 4.66 bcd	2.29 ± 0.12 ab	256.75 ± 16.49 b	37 ± 5.73 a
A5-3	83.37 ± 2.09 cde	2.02 ± 0.16 abc	225.25 ± 28.20 bcd	36 ± 4.46 ab
A8-3	83.12 ± 3.33 cde	1.99 ± 0.14 abc	188.75 ± 3.25 de	29.5 ± 1.22 abcd
A9-3	81.75 ± 4.07 def	1.67 ± 0.08 c	134.62 ± 7.03 f	18.375 ± 1.03 def
A10-3	80.62 ± 1.64 ef	2.04 ± 0.06 abc	210.12 ± 7.72 cde	31.25 ± 2.62 abc
A1-4	87.12 ± 1.76 bcde	1.79 ± 0.1 bc	233.12 ± 10.99 bc	37.25 ± 4.25 a
A2-4	83.75 ± 2.25 cde	1.71 ± 0.06 c	184.5 ± 4.32 e	25.75 ± 1.37 bcdef
A3-4	91.37 ± 1.37 bc	2.30 ± 0.08 ab	186.12 ± 7.75 e	28.75 ± 3.62 abcde
A4-4	103.5 ± 9.67 a	1.99 ± 0.11 abc	228.12 ± 16.32 bc	39.25 ± 5.99 a
A5-4	83.5 ± 2.46 cde	1.87 ± 0.16 bc	219.5 ± 8.87 cde	39.25 ± 6.12 a
A6-4	84.87 ± 2.56 cde	1.98 ± 0.04 abc	199.5 ± 2.84 cde	33.5 ± 2.08 abc
A7-4	55.12 ± 1.53 h	1.75 ± 0.19 c	145.25 ± 4.63 f	24.12 ± 1.8 cdef
A8-4	69 ± 2.07 g	1.8 ± 0.03 bc	148.87 ± 3.39 f	24 ± 1.99 cdef
A10-4	73.5 ± 2.16 fg	1.93 ± 0.11 bc	141.37 ± 5.25 f	16.87 ± 2.02 f

UC: Uninoculated and untreated control. Results are presented as means ± SE ($n = 10$, $P \leq 0.05$). For each column, values followed by the same letter are not significantly different according to Duncan's Multiple Range test (at $P \leq 0.05$).

4. Discussion

Several investigations have reported that microorganisms isolated from composts could enhance the growth of various crops and controlling several phytopathogens (Coelho et al., 2020; Joshi et al., 2009). In previous work,

we demonstrated the capacity of composts and compost teas to suppress stem rot disease caused by *S. rolfsii* and to promote tomato growth (Ayed et al., 2018b; 2018c). Therefore, this study was more focused on the isolation and screening of actinomycetes naturally associated with 2 composts as biocontrol and

growth-promoting agents. A total of 17 actinomycetes isolates were recovered from 2 selected composts C₃ and C₄. When assessed for their ability to suppress stem rot disease, 12 actinobacterial isolates (namely A2-3, A3-3, A4-3, A5-3, A8-3, A9-3, A1-4, A2-4, A3-4, A4-4, A6-4, and A10-4) had successfully decreased disease severity on *S. rolf sii*-inoculated tomato cv. Rio Grande seedlings. The noted antifungal activity of actinomycetes isolates confirmed the results obtained by Anusree and Bhai (2017) who demonstrated their ability to protect pepper by 98.10% from *Sclerotium* foot rot disease. Also, Errakhi et al. (2007) reported the antagonistic potential of actinobacterial isolates against *S. rolf sii* infecting sugar beet. Several other investigations indicated that actinomycetes are promising microbial agents for the biological control of numerous fungal and bacterial plant pathogens (Dandan et al., 2018; Mingma et al., 2014; Golinska & Dahm, 2013; Kobayashi et al., 2012; Patil et al., 2010). Previous studies have also shown that actinomycetes can produce plant growth promoters (Srivastava et al., 2015; Gopalakrishnan et al., 2014). In the current study, some compost-associated actinomycetes were screened for their ability to promote the growth of tomato seedlings. Interestingly, our results demonstrated that A3-3, A4-3, A2-4, A3-4, and A4-4 isolates exhibited the highest growth-promoting potentials on *S. rolf sii*-inoculated tomato seedlings and that A3-3 and A4-4 were also the most effective on pathogen-free seedlings. Nevertheless, A1-3, A7-4, A8-4, and A10-4 isolates did not exhibit any positive effect on seedling growth. Thus, actinomycetes associated with composts can be explored as potential agents for seedling growth-stimulation and stem rot disease biocontrol. These effects may be explained by the ability of plant root exudates to stimulate the growth

of actinomycetes and the last ones use these exudates for the synthesis of antimicrobial substances active against plant pathogens (Yuan & Crawford, 1995). Most actinomycetes belong to the genus *Streptomyces* and are well known as the main sources of bioactive secondary metabolites (Khanna et al., 2011; Goodfellow & Simpson 1987). Therefore, different aspects of these microorganisms have been studied, such as the production of metabolites that control plant disease and/or improve plant growth. To inhibit phytopathogens, the microbial biocontrol agents use different mechanisms such as the production of cell wall-degrading enzymes, parasitism, antibiosis, and the induction of host resistance (Palaniyandi et al., 2013). In this regard, previous findings showed the ability of actinomycetes to synthesize hydrolytic enzymes including chitinases, proteases, amylases, lipases, cellulases, or β -1,3 glucanase (Anusree & Bhai, 2017; Jayamurthy et al., 2014). In the same sense, actinomycetes are described as natural antibiotic producers and can inhibit various soilborne pathogens such as *Fusarium oxysporum* f. sp. *cubense* (Getha & Vikineswary, 2002), *Verticillium dahliae* (Aouar et al., 2012), and *Rhizoctonia solani* (Patil et al., 2010). Regarding plant growth promotion, the efficiency of various microorganisms, including actinomycetes, were widely reported (Himaman et al., 2016; Gopalakrishnan et al., 2014; Salla et al., 2014; Peralta et al., 2012). These agents may promote their host growth either directly by the synthesis of phytohormones (Goudjal et al., 2013), nitrogen fixation (Bibha et al., 2017), siderophores, phosphate and zinc solubilization, indole acetic acid, extracellular enzyme lipase (Anusree & Bhai, 2017); or indirectly through the suppression of their associated plant pathogens (Jose & Jha, 2016).

5. Conclusion

Screened composts were found to be a potentially important source for the isolation of potent actinomycetes, acting both as biocontrol agents and as biofertilizers. Nevertheless, more investigations are needed to more understand the mechanisms of action, to identify secondary metabolites involved in the growth promotion and the stem rot biocontrol, and to identify the potent isolates; which can be used for the development of eco-friendly and economically feasible biofertilizers and bio-fungicides.

Acknowledgments

This work was funded by the Ministry of Higher Education and Scientific Research in Tunisia through the budget assigned to UR13AGR09-Integrated Horticultural Production in the Tunisian Centre-East, The Regional Research Centre on Horticulture and Organic Agriculture of Chott-Meriem, University of Sousse, Tunisia.

References

- Alfano G, Lustrato G, Lima G, Vitullo D, Ranalli G, 2011. Characterization of composted olive mill wastes to predict potential plant disease suppressiveness. *Biological Control* **58**(3): 99–207.
- Aouar L, Lerat S, Ouffroukh A, Boulahrouf A, Beaulieu C, 2012. Taxonomic identification of rhizospheric actinobacteria isolated from Algerian semi-arid soil exhibiting antagonistic activities against plant fungal pathogens. *Canadian Journal of Plant Pathology* **34**: 165–176.
- Anusree T, Bhai RS, 2017. Rhizosphere Actinobacteria for combating *Phytophthora capsici* and *Sclerotium rolfsii*, the major soil borne pathogens of black pepper (*Piper nigrum* L). *Biological Control* **109**: 1–13.
- Aouar L, Lerat S, Ouffroukh A, Boulahrouf A, Beaulieu C, 2012. Taxonomic identification of rhizospheric actinobacteria isolated from Algerian semi-arid soil exhibiting antagonistic activities against plant fungal pathogens. *Canadian Journal of Plant Pathology* **34**(2): 165–176.
- Aycock R, 1966. Stem rot and other diseases caused by *Sclerotium rolfsii* or the status of Rolf's fungus after 70 years. *Technical Bulletin/North Carolina Agricultural Experiment Station* **174**: 136–202.
- Aydi-Ben Abdallah R, Jabnoun-Khiareddine H, Nefzi A, Ayed F, Daami-Remadi M, 2018. Endophytic bacteria from *Solanum nigrum* with plant growth-promoting and Fusarium wilt-suppressive abilities in tomato. *Tunisian Journal of Plant Protection* **13**(2): 157–182.
- Aydi-Ben Abdallah R, Jabnoun-Khiareddine H, Daami-Remadi M, 2020. Variation in the composition of potato plants depending on cropping season, cultivar type, and plant development stage. *International Journal of Agriculture, Environment and Food Science* **4**(3): 319–333.
- Ayed F, Jabnoun-Khiareddine H, Aydi-Ben-Abdallah R, Daami-Remadi M, 2018a. Effect of temperatures and culture media on *Sclerotium rolfsii* mycelial growth, sclerotial formation and germination. *Journal of Plant Pathology and Microbiology* **09**: 429.
- Ayed F., Jabnoun-Khiareddine H. and Daami-Remadi M, 2018b. Valorisation du compost comme moyen de lutte contre *Sclerotium rolfsii* agent responsable de la

- pourriture des tiges. Actes des 9^{èmes} Journées Internationales sur la Valorisation des Bioressources à Monastir, Tunisia, C.O. 48.
- Ayed F., Jabnoun-Khiareddine H., Aydi Ben Abdallah R. and Daami-Remadi M, 2018c. Sclerotium collar rot suppression and tomato growth promotion achieved using different compost teas. Proceedings of the fifth International Conference on Sustainable Agriculture and Environment, Hammamet, Tunisia, pp. 61.
- Bhatti AA, Haq S, Bhat RA, 2017. Actinomycetes benefaction role in soil and plant health. *Microbial Pathogenesis* **111**: 458–467.
- Bibha D, Githanjali N, Lora P, Volker SB, 2017. Diversity of free living nitrogen fixing *Streptomyces* in soils of the badlands of South Dakota. *Microbiological Research* **195**: 31–39.
- Bohacz J, 2018. Microbial strategies and biochemical activity during lignocellulosic waste composting in relation to the occurring biothermal phases. *Journal of Environmental Management* **206**: 1052–1062.
- Coelho L, Reis M, Guerrero C, Dionísio L, 2020. Use of organic composts to suppress bentgrass diseases in *Agrostis stolonifera*. *Biological control* **141**: 104154.
- Coventry E, Noble R, Mead A, Marin F, Perez J, Whipps J, 2006. Allium white rot suppression with composts and *Trichoderma viridae* in relation to sclerotia viability. *Phytopathology* **96**(9): 1009–1020.
- Daami-Remadi M, Jabnoun-Khiareddine H, Sdiri A, El Mahjoub M, 2012. Comparative reaction of potato cultivars to *Sclerotium rolfii* assessed by stem rot and tuber decay severity. *Pest Technology* **6**(Special Issue 1): 54–59.
- Dandan H, Lanying W, Yanping L, 2018. Isolation, identification, and the growth promoting effects of two antagonistic actinomycete strains from the rhizosphere of *Mikania micrantha* Kunth. *Microbiological Research* **208**: 1–11.
- De Corato U, 2020. Disease-suppressive compost enhances natural soil suppressiveness against soil-borne plant pathogens: A critical review. *Rhizosphere* **130**: 100192.
- De Curtis F, Lima G, Vitullo D, De Cicco V, 2010. Biocontrol of *Rhizoctonia solani* and *Sclerotium rolfii* on tomato by delivering antagonistic bacteria through a drip irrigation system. *Crop Protection* **29**: 663–670.
- Errakhi R, Bouteau F, Lebrihi A, Barakate M, 2007. Evidences of biological control capacities of *Streptomyces* spp. against *Sclerotium rolfii* responsible for damping-off disease in sugar beet (*Beta vulgaris* L.). *World Journal of Microbiology and Biotechnology* **23**: 1503–1509.
- Fery RL, Dukes PD, 2002. Southern Blight (*Sclerotium rolfii* Sacc.) of Cowpea: Yield-loss estimates and sources of resistance. *Crop Protection* **21**: 403–408.
- Getha K, Vikineswary S, 2002. Antagonistic effects of *Streptomyces violaceusniger* strain G10 on *Fusarium oxysporum* f.sp. *cubense* race 4: indirect evidence for the role of antibiosis in the antagonistic process. *Journal of Industrial Microbiology & Biotechnology* **28**(6): 303–310.
- Golinska P, Dahm H, 2013. Antagonistic properties of *Streptomyces* isolated from forest soils against fungal pathogens of pine seedlings. *Dendrobiology* **69**: 87–

- 97.
- Goodfellow M, Simpson KE, 1987. Ecology of Streptomycetes. *Frontiers in Applied Microbiology* **2**: 97–125.
- Gopalakrishnan S, Vadlamudi S, Bandikinda P, Sathya A, Vijayabharathi R, Rupela O, Kudapa H, Katta K, Varshney RK, 2014. Evaluation of *Streptomyces* strains isolated from herbal vermicompost for their plant growth-promotion traits in rice. *Microbiological Research* **169**: 40–48.
- Goudjal Y, Toumatia O, Sabaou N, Barakate M, Mathieu F, Zitouni A, 2013. Endophytic actinomycetes from spontaneous plants of Algerian Sahara: indole-3-acetic acid production and tomato plants growth promoting activity. *World Journal of Microbiology and Biotechnology* **29**: 1821–1829.
- Hadar Y, Papadopoulou KK, 2012. Suppressive composts: microbial ecology links between abiotic -environments and healthy plants. *Annual Review of Phytopathology* **50**: 133–153.
- Hamdali H, Hafidi M, Virolle MJ, Ouhdouch Y, 2008. Rock phosphate-solubilizing Actinomycetes: screening for plant growth-promoting activities. *World Journal of Microbiology and Biotechnology* **24**: 2565–2575.
- Himaman W, Thamchaipenet A, Pathomaree W, Duangmal K, 2016. Actinomycetes from Eucalyptus and their biological activities for controlling Eucalyptus leaf and shoot blight. *Microbiological Research* **188-189**: 42–52.
- Jayamurthy H, Valappil K, Dastagar SG, Pandey A, 2014. Anti-fungal potentials of extracellular metabolites of Western Ghats isolated *Streptomyces* sp. NII 1006 against moulds and yeasts. *Indian Journal of Experimental Biology* **52**: 1138–1146.
- Jose PA, Jha B, 2016. New dimensions of research on Actinomycetes: quest for next generation antibiotics. *Frontiers in Microbiology* **7**: 1295.
- Joshi D, Hooda KS, Bhatt JC, Mina BL, Gupta HS, 2009. Suppressive effects of composts on soil-borne and foliar diseases of French bean in the field in the western Indian Himalayas. *Crop Protection* **28**: 608–615.
- Kator L, Hosea ZY, Oche OD, 2015. *Sclerotium rolfsii*; Causative organism of Southern blight, stem rot, white mold and sclerotia rot disease. *Annals of Biological Research* **6**: 78–89.
- Khanna M, Solanki R, Lal R, 2011. Selective isolation of rare actinomycetes producing novel antimicrobial compounds. *International Journal of Advanced Biotechnology and Research* **2**: 357–375.
- Kobayashi YO, Kobayashi A, Maeda M, Takenaka S, 2012. Isolation of antagonistic *Streptomyces* sp.: against a potato scab pathogen from a field cultivated with wild oat. *Journal of General Plant Pathology* **78**: 62–72.
- Larkin RP, Tavantzis S, 2013. Use of biocontrol organisms and compost amendments for improved control of soilborne diseases and increased potato production. *American Journal of Potato Research* **90**: 261–270.
- Manici LM, Caputo F, Babini V, 2004. Effect of green manure on *Pythium* sp. population and microbial communities in intensive cropping systems. *Plant Soil* **263**: 133–142.
- Martin SCCG, 2015. Enhancing soil suppressiveness using compost and compost tea. In: Meghvansi MK, Varma A, eds. *Organic amendments and soil suppressiveness in plant disease management*. *Soil Biology*, Volume 46,

- Springer International Publishing, Switzerland, pp. 25–49.
- Mingma R, Pathom-aree W, Trakulnaleamsai S, Thamchaipenet A, Duangmal K, 2014. Isolation of rhizospheric and roots endophytic actinomycetes from *Leguminosae* plant and their activities to inhibit soybean pathogen *Xanthomonas campestris* pv. *glycine*. *World Journal of Microbiology & Biotechnology* **30**: 271–280.
- Nurkanto A, Julistiono H, 2014. Screening and study of antifungal activity of leaf litter actinomycetes isolated from Ternate island, Indonesia. *Asian Pacific Journal of Tropical Medicine* **7**: S238–S243.
- Palaniyandi SA, Yang SH, Zhang L, Suh JW, 2013. Effects of actinobacteria on plant disease suppression and growth promotion. *Applied Microbiology and Biotechnology* **97**: 9621–9636.
- Palla MS, Guntuku GS, Muthyala MKK, Pingali S, Sahu PS, 2018. Isolation and molecular characterization of antifungal metabolite producing actinomycete from mangrove soil. *Beni-Suef University Journal of Basic and Applied Sciences* **7**: 250–256.
- Pane C, Piccolo A, Spaccini R, Celano G, Vilecco D, Zaccardelli M, 2013. Agricultural waste-based composts exhibiting suppressivity to diseases caused by the phytopathogenic soil-borne fungi *Rhizoctonia solani* and *Sclerotinia minor*. *Applied Soil Ecology* **65**: 43–51.
- Pane C, Spaccini R, Piccolo A, Celano G, Zaccardelli M, 2019. Disease suppressiveness of agricultural green-waste composts as related to chemical and bio-based properties shaped by different on-farm composting methods. *Biological Control* **137**: 104026.
- Patil HJ, Srivastava AK, Kumar S, Chaudhari BL, Arora DK, 2010. Selective isolation, evaluation and characterization of antagonistic actinomycetes against *Rhizoctonia solani*. *World Journal of Microbiology and Biotechnology* **26**(12): 2163–2170.
- Peralta KD, Araya T, Valenzuela S, Sossa K, Martínez M, Cortés HP, Sanfuentes E, 2012. Production of phytohormones, siderophores and population fluctuation of two root-promoting rhizobacteria in *Eucalyptus globulus* cuttings. *World Journal of Microbiology and Biotechnology* **28**: 2003–2014.
- Punja ZK, 1985. The biology, ecology and control of *Sclerotium rolfsii*. *Annual Review of Phytopathology* **23**: 97–127.
- Qin S, Xing K, Jiang JH, Xu LH, Li WJ, 2011. Biodiversity, bioactive natural products and biotechnological potential of plant-associated endophytic actinobacteria. *Applied Microbiology and Biotechnology* **89**: 457–473.
- Salla TD, da Silva TR, Astarita LV, Santarém ER, 2014. *Streptomyces* rhizobacteria modulate the secondary metabolism of *Eucalyptus* plants. *Plant Physiology and Biochemistry* **85**: 14–20.
- Scotti R, Pane C, Spaccini R, Palese AM, Piccolo A, Celano G, Zaccardelli M, 2016. On-farm compost: a useful tool to improve soil quality under intensive farming systems. *Applied Soil Ecology* **107**: 13–23.
- Singh SP, Singh HB, Singh DK, 2013. *Trichoderma harzianum* and *Pseudomonas* sp. mediated management of *Sclerotium rolfsii* rot in tomato (*Lycopersicon esculentum* Mill.). *Life Sciences* **8**: 801–804.
- Sridharan AP, Sugitha T, Karthikeyan G, Sivakumar U, 2020. Comprehensive profiling of the VOCs of *Trichoderma*

- longibrachiatum* EF5 while interacting with *Sclerotium rolfii* and *Macrophomina phaseolina*. *Microbiological Research* **236**: 126436.
- Srivastava S, Patel JS, Singh HB, Sinha A, Sarma BK, 2015. *Streptomyces rochei* SM3 induces stress tolerance in chickpea against *Sclerotinia sclerotiorum* and NaCl. *Journal of Phytopathology* **163**: 583–592.
- Stavi I, Bel G, Zaady E, 2016. Soil functions and ecosystem services in conventional, conservation, and integrated agricultural systems. A review. *Agronomy for Sustainable Development* **36**: 32.
- Sun S, Sun F, Deng D, Zhu X, Duan C, Zhu Z, 2020. First report of southern blight of mung bean caused by *Sclerotium rolfii* in China. *Crop Protection* **130**: 105055.
- Tubeileh AM, Stephenson GT, 2020. Soil amendment by composted plant wastes reduces the *Verticillium dahliae* abundance and changes soil chemical properties in a bell pepper cropping system. *Current Plant Biology* **22**: 100148.
- Yuan WM, Crawford DL, 1995. Characterization of *Streptomyces lydicus* WYEC108 as a potential biocontrol agent against fungal root and seed rots. *Applied and Environmental Microbiology* **61**: 3119–3128.